


Usual source(s) for Sick and Well Care
 Percent of CSHCN who have usual source(s) for sick and well care
2009/10 National Survey of Children with Special Health Care Needs



Usual source(s) for Sick and Well Care

Percent of CSHCN who have usual source(s) for sick and well care

2009/10 National Survey of Children with Special Health Care Needs

US Prevalence		89.3%
HIGHER than US; Statistically Significant	Connecticut	93.9%
	Alabama	93.4%
	Kentucky	92.6%
	New Hampshire	92.5%
	Iowa	92.2%
	West Virginia	92.1%
	Maryland	92.0%
	Oregon	91.9%
	Pennsylvania	91.9%
	North Carolina	91.8%
HIGHER than US; Not Statistically Significant	Massachusetts	91.6%
	New York	91.6%
	Virginia	91.5%
	Hawaii	91.2%
	Delaware	91.2%
	Ohio	91.0%
	Maine	90.9%
	Kansas	90.7%
	Rhode Island	90.7%
	Nebraska	90.5%
	Michigan	90.3%
	Minnesota	90.3%
	South Carolina	90.3%
	Washington	90.1%
District of Columbia	90.0%	
Oklahoma	89.4%	
LOWER than US; Not Statistically Significant	California	89.3%
	Indiana	89.2%
	Georgia	88.9%
	Vermont	88.8%
	Colorado	88.7%
	New Jersey	88.4%
	Wisconsin	88.4%
	Montana	88.3%
	Tennessee	88.1%
	Florida	87.9%
	Alaska	87.7%
	Arkansas	87.6%
	Utah	87.5%
	Missouri	87.4%
	Louisiana	87.1%
	New Mexico	87.1%
	North Dakota	86.5%
	Wyoming	86.5%
	Nevada	86.2%
Idaho	85.8%	
South Dakota*	83.3%	
LOWER than US; Stat Sig	Texas*	86.1%
	Mississippi	85.7%
	Illinois	85.2%
	Arizona	84.0%

NOTES:

Higher %'s = Better Performance

Statistical Significance: $p < .05$