


**Indicator 8: CSHCN needing a referral for specialist care or services and had problems getting it**  
 Percent of CSHCN who met criteria for Indicator #8  
**2009/10 National Survey of Children with Special Health Care Needs**



## Indicator 8: CSHCN needing a referral for specialist care or services and had problems getting it

Percent of CSHCN who needed referrals who met criteria for Indicator #8

### 2009/10 National Survey of Children with Special Health Care Needs

US Prevalence		23.4%
LOWER than US; Statistically Significant	Alabama	12.6%
	Tennessee	13.5%
	Indiana	13.7%
	Kansas	14.0%
	New Hampshire	14.0%
	Vermont	15.1%
	Minnesota	15.6%
	Massachusetts	15.7%
	Nebraska	15.8%
	North Carolina	16.0%
	Maine	16.1%
	Pennsylvania	17.2%
LOWER than US; Not Significant	Georgia	17.3%
	Mississippi	17.8%
	Kentucky	18.8%
	Iowa	19.1%
	Delaware	19.1%
	Missouri	19.2%
	Rhode Island	20.1%
	Idaho	20.2%
	Washington	20.9%
	West Virginia	21.1%
	District of Columbia	21.1%
	Ohio	21.1%
	Virginia	21.2%
	North Dakota	21.8%
	Illinois	22.4%
	Wisconsin	22.5%
Hawaii	22.7%	
Alaska	22.7%	
HIGHER than US; Not Significant	Oklahoma	23.4%
	New Jersey	23.8%
	Maryland	23.8%
	Oregon	23.9%
	Arkansas	24.0%
	Florida	24.1%
	Montana	24.4%
	Wyoming	24.8%
	New York	25.2%
	Connecticut	25.4%
	New Mexico	25.7%
	South Dakota	25.8%
	Louisiana	26.1%
	South Carolina	26.8%
	Michigan	28.0%
	Utah	28.6%
	Arizona	30.2%
Colorado	30.5%	
HIGHER than US; SS	Texas	31.3%
	California	33.9%
	Nevada	35.8%

#### NOTES:

Lower %'s = Better performance

Statistical significance:  $p < .05$

\* Due to sampling error and other factors, some state estimates differ statistically from the national prevalence while other state estimates of similar or equal size do not.