


Medical Home Component – Effective Care Coordination

Percent of children who received effective care coordination when needed

2005/06 National Survey of Children with Special Health Care Needs



Medical Home Component – Care Coordination

Percent of children who received effective care coordination when needed

2005/06 National Survey of Children with Special Health Care Needs

US Prevalence		59.2%
HIGHER than US; Stat. Sig.	Kansas	68.0%
	Ohio	67.3%
	Iowa	66.8%
	Nebraska	64.9%
	Louisiana	64.6%
	Missouri	64.1%
HIGHER than US; Not Significant	Kentucky	63.4%
	South Carolina	63.4%
	South Dakota	63.3%
	Indiana	63.2%
	Vermont	63.1%
	North Dakota	63.0%
	Wisconsin	63.0%
	Tennessee	62.9%
	Montana	62.8%
	West Virginia	62.8%
	Rhode Island	62.5%
	Maine	62.5%
	Alabama	62.2%
	Arkansas	61.9%
	Michigan	61.4%
	Mississippi	61.1%
	Minnesota	61.0%
	Hawaii	60.9%
	Pennsylvania	60.4%
	Texas	60.4%
Colorado	60.0%	
Georgia	59.5%	
North Carolina	59.2%	
LOWER than US; Not Significant	Oklahoma	59.1%
	Wyoming	58.6%
	New York	58.5%
	New Hampshire	58.4%
	Maryland	57.6%
	Connecticut	57.6%
	Illinois	57.6%
	Virginia	56.8%
	New Mexico	56.6%
	Oregon	56.3%
	Idaho	55.8%
	Delaware	55.5%
	Utah	55.4%
	Florida	55.3%
	Massachusetts	55.0%
	California	54.5%
LOWER than US; Stat. Sig.	Washington	54.2%
	Arizona	52.9%
	Alaska	52.7%
	District of Columbia	51.1%
	New Jersey	51.0%
	Nevada	50.6%

NOTES:

Higher %'s = Better Performance

Statistical Significance: $p < .05$