

Medical Home Component – Usual Source for Sick and Well Care

Percent of children with a usual source for sick and well care

2005/06 National Survey of Children with Special Health Care Needs

US Prevalence		92.9%
HIGHER than US; Stat. Sig.	New Hampshire	95.9%
	Washington	95.6%
	Iowa	95.4%
	Nebraska	95.2%
	Connecticut	95.1%
	Pennsylvania	95.1%
	Utah	95.0%
HIGHER than US; Not Significant	Maine	94.6%
	Minnesota	94.6%
	Rhode Island	94.5%
	Delaware	94.5%
	Massachusetts	94.4%
	Wisconsin	94.4%
	Illinois	94.1%
	Ohio	94.0%
	South Dakota	94.0%
	Kansas	93.9%
	New Jersey	93.9%
	Vermont	93.8%
	Maryland	93.7%
	Virginia	93.6%
	South Carolina	93.2%
	Kentucky	93.2%
	Tennessee	93.0%
North Carolina	93.0%	
LOWER than US; Not Significant	West Virginia	92.9%
	Florida	92.9%
	Wyoming	92.9%
	Texas	92.9%
	Indiana	92.8%
	Michigan	92.7%
	New Mexico	92.6%
	Oklahoma	92.5%
	Alabama	92.5%
	Idaho	92.4%
	Oregon	92.4%
	New York	92.3%
	Georgia	92.2%
	Hawaii	91.9%
	Arkansas	91.7%
	Colorado	91.6%
	Missouri	91.4%
	Alaska	91.2%
	California	91.1%
	North Dakota	90.9%
Mississippi	90.9%	
District of Columbia	90.5%	
LOWER than US; Stat. Sig.	Montana	90.2%
	Arizona	89.9%
	Nevada	89.4%
	Louisiana	88.3%

NOTES:

Higher %'s = Better Performance

Statistical Significance: $p < .05$